

CHAPTER ELEVEN

DIAGNOSIS AND TREATMENT OF ILLNESS

Dehydration and Fluid Replacement Therapy

The average daily water budget for a bat is determined by what they take in (food and water), and what is lost in feces and urine, as well as evaporative loss that occurs during flight and normal activities. The average daily water budget is 24.5% and 15.8% of the body weight for a big brown bat (*E. fuscus*), and Brazilian free-tailed bat (*T. brasiliensis*) respectively. Without sufficient intake, a 10 to 20g bat can have a daily water loss of as much as 1 to 3mls.



Figure 11-1. Glossy wings on a well-hydrated bat. *E. fuscus*. Bat World facility. Photo by A. Lollar.

Bats that have been confined in areas without food or water, or are too weak to fly due to injury or illness will become dehydrated. Bats in captivity can become dehydrated if the humidity is too low or they don't have access to fresh water at all times. Orphaned bats are always dehydrated to some degree.

Well hydrated bats will have smooth and slippery mucous membranes and the wing membrane will appear glossy (Figure 11-1). A bat that is slightly dehydrated (around 5%) will have slightly dry mucous membranes and may have stringy saliva. Other signs include loss of appetite, vomiting, and feces stuck to the tail membrane. The skin of a bat that is over 5% dehydrated will take longer than one second to return to normal after being tented, and the mucus membranes will appear dry and tacky. A bat that is around 8% dehydrated will have a flattened or sunken abdomen with wrinkled skin, and the wing membrane may appear somewhat dull. The eyes of a severely dehydrated bat (over 10%) will actually sink back in the sockets (and the bat may be unable to open its eyes). The skin will remain in place when pinched. Inability to coordinate muscular movements may be observed, or the bat may be comatose. Bats over 10% dehydrated may not survive.

Dehydrated bats cannot be rehydrated with oral fluids alone and therefore must be given warmed subcutaneous (SQ) injections of electrolytes (e.g., lactated Ringers solution, Normasol or Plasmalyte), regardless of whether or not they accept fluids orally. Depending on the severity of dehydration, electrolytes administered SQ are usually absorbed within 5-30 minutes (see Injection Technique on the following page). Severely dehydrated bats that do not absorb fluids may be suffering from irreparable organ failure.

After an initial injection the skin of a bat quickly resumes its former elasticity. Most adult bats will also be sufficiently hydrated after only 1 or 2 injections. Once hydrated, infant bats can be fed (see Feeding and Care of Infant Bats) and adult bats can be offered small amounts of soft food (see Feeding Adult Bats). The Bat World Sanctuary soft food diet is easier to swallow, has complete nutrition, and contains additional fluids which will assist in the rehydration process. Note: Signs of over-hydration include shivering and edema around the face,

Table 11-1
Fluid Replacement Schedule

Weight (g)	Dose (ml)	Frequency (hours)
Pups:		
Less than 1.0	0.50	1 injection every 24 hours
1.0 to 5.0	0.75	1 to 2 injections within 24 hours
Juveniles or Adults:		
3.0 to 5.0	1.0	1 to 2 injections within 24 hours
5.1 to 15.0	1.0 to 3.0	1 to 2 injections within 24 hours
15.1 to 20.0	2.0 to 3.0	1 to 2 injections within 24 hours
20.0 to 65.0	5.0 to 15.0	1 to 2 injections within 24 hours